

RESEARCH PAPER RP1525

*Part of Journal of Research of the National Bureau of Standards, Volume 30,
February 1943*

MINIATURE GEIGER-MÜLLER COUNTER

By Leon F. Curtiss

ABSTRACT

A miniature Geiger-Müller tube counter having a tube 0.8 mm in internal diameter is described. This "hypodermic-needle" type of counter has been found useful in locating and measuring a well-defined beam of gamma radiation. It could also be used to measure a beam of hard X-rays reflected from a crystal. Another suggested use is direct insertion into specimens, such as living biological material.

There are many applications for which a Geiger-Müller counter of very small size would be useful. Where it is desirable to measure and locate a narrow beam of radiation, it would be very convenient to have a counter approximately the size of the cross section of the beam. This requires a counter of the order of 1 mm in diameter. The author knows of no previous attempts to construct counters of such size.

The ease with which thoroughly reliable counters containing alcohol vapor and argon may be constructed led to the trial of small counters. No difficulty was encountered in constructing counters with an internal diameter of 3 mm. Counters of this size give slightly larger pulses than is obtained from counters 2 or 3 cm in diameter. They also show a reasonable plateau and are quite stable. Counters of this diameter would not serve to locate a well-defined beam of radiation with sufficient precision for some purposes.

Encouraged by the results with the 3-mm counters, the author constructed the counter shown in figure 2. The No. 2 sewing needle is included in the picture for size comparison. The tube of the counter has an internal diameter of 0.8 mm and a length of about 3 mm. The central wire is tungsten, 0.05 mm in diameter. The filling gas is a mixture of amyl acetate vapor and argon at about 4-cm pressure.

This counter has an extremely low blank, averaging around 20 pulses per hour. The pulses are even larger than those from the 3-mm counters. This increase of pulse size with decrease of tube diameter could be expected from the known fact that the ionization is confined to a thin sheath near the central wire. The thickness of this sheath is increased as the tube becomes smaller, since the ratio of maximum to minimum gradient is smaller for the counter tube of smaller diameter.

The plateau for this counter is shown in figure 1. Although it is considerably steeper than for larger counters, this counter yields good results when operated from a stabilized voltage. This particular

counter has been in use for several months with no sign of deterioration.

The shape of the counter resembles a large hypodermic needle, which suggests another type of use. It could be introduced directly into material to be tested and therefore increase the sensitivity of the measurement. This could be of value in studying biological specimens. It could even be used with live animals. No attempts have been made to construct counters of smaller diameter, but it seems likely they could be made. The chief difficulty is in devising a method of locating the central wire so that it does not touch the counter tube. Exact centering of this wire is not necessary, since the only precaution taken with the present counter was to make certain that the wire was not in contact with the tube.




FIGURE 1.—*Hypodermic-needle-type Geiger-Müller counter.*

Further work is in progress in an effort to improve the characteristics of these miniature tube counters and to determine the lowest limit of tube diameter that can be used.

WASHINGTON, December 2, 1942.

○




FIGURE 2.—*Miniature Geiger-Müller counter.*

The sewing needle is included for size comparison.

NATIONAL BUREAU OF STANDARDS,
WASHINGTON, D. C.

Send me the Mathematical Tables marked X below. I enclose remittance ¹ to cover the cost.

Mark X	Title of publication	Price		Amount enclosed
		United States and its possessions, and countries ex- tending franking privilege	Other countries	
-----	MT1. Table of the first ten powers of the integers from 1 to 1000-----	\$0. 50	\$0. 65	-----
-----	MT2. Tables of the exponential function e^x -----	2. 00	2. 50	-----
-----	MT3. Tables of circular and hyperbolic sines and cosines for radian arguments--	2. 00	2. 50	-----
-----	MT4. Tables of sines and cosines for radian arguments-----	2. 00	2. 50	-----
-----	MT5. Tables of sine, cosine, and exponential integrals, volume I-----	2. 00	2. 50	-----
-----	MT6. Tables of sine, cosine, and exponential integrals, volume II-----	2. 00	2. 50	-----
-----	MT7. Table of natural logarithms, volume I-----	2. 00	2. 50	-----
-----	MT8. Tables of probability functions, volume I-----	2. 00	2. 50	-----
-----	MT9. Table of natural logarithms, volume II-----	2. 00	2. 50	-----
-----	MT10. Table of natural logarithms, volume III-----	2. 00	2. 50	-----
-----	MT11. Tables of moments of inertia and section moduli-----	2. 00	2. 50	-----
-----	MT12. Table of natural logarithms, volume IV-----	2. 00	2. 50	-----
-----	MT13. Table of sine and cosine integrals for arguments from 10 to 100-----	2. 00	2. 50	-----
-----	MT14. Tables of probability functions, volume II-----	2. 00	2. 50	-----
-----	MT15. The hypergeometric and Legendre functions with applications to integral equations of potential theory-----	2. 00	2. 50	-----
	Total remittance-----			

¹ Remittance should be in form of post-office money order, or check, and made payable to the order of the "National Bureau of Standards" in United States currency.

Sent to-----

Number and Street-----

City and State-----

(Cut here)

MATHEMATICAL TABLES

Attention is invited to a series of publications which is being prepared by the *Project for the Computation of Mathematical Tables* conducted by the Federal Works Agency, Work Projects Administration for the City of New York under the sponsorship of the National Bureau of Standards. The tables which have been made available through the National Bureau of Standards are listed below.

There is included in this list a publication on the hypergeometric and Legendre functions (MT15), prepared by the Bureau.

MT1. TABLE OF THE FIRST TEN POWERS OF THE INTEGERS FROM 1 TO 1000:

(1938) VIII+80 pages; heavy paper cover. 50 cents.

MT2. TABLES OF THE EXPONENTIAL FUNCTION e^x :

The ranges and intervals of the argument and the number of decimal places in the entries are given below:

Range of x	Interval of x	Decimals given
-2.5000 to 1.0000	0.0001	18
1.0000 to 2.5000	.0001	15
2.500 to 5.000	.001	15
5.00 to 10.0	.01	12

(1939) XV+535 pages; bound in buckram, \$2.00.

MT3. TABLES OF CIRCULAR AND HYPERBOLIC SINES AND COSINES FOR RADIAN ARGUMENTS:

Contains 9 decimal place values of $\sin x$, $\cos x$, $\sinh x$ and $\cosh x$ for x (in radians) ranging from 0 to 2 at intervals of 0.0001.

(1939) XVII+405 pages; bound in buckram, \$2.00.

MT4. TABLES OF SINES AND COSINES FOR RADIAN ARGUMENTS:

Contains 8 decimal place values of sines and cosines for radian arguments ranging from 0 to 25 at intervals of 0.001.

(1940) XXIX+275 pages; bound in buckram, \$2.00.

MT5. TABLES OF SINE, COSINE, AND EXPONENTIAL INTEGRALS, VOLUME I:

Values of these functions to 9 places of decimals from 0 to 2 at intervals of 0.0001.

(1940) XXVI+444 pages; bound in buckram, \$2.00.

MT6. TABLES OF SINE, COSINE, AND EXPONENTIAL INTEGRALS, VOLUME II:

Values of these functions to 9, 10, or 11 significant figures from 0 to 10 at intervals of 0.001 with auxiliary tables.

(1940) XXXVII+225 pages; bound in buckram, \$2.00.

MT7. TABLE OF NATURAL LOGARITHMS, VOLUME I:

Logarithms of the integers from 1 to 50,000 to 16 places of decimals.

(1941) XVIII+501 pages; bound in buckram, \$2.00.

MT8. TABLES OF PROBABILITY FUNCTIONS, VOLUME I:

Values of these functions to 15 places of decimals from 0 to 1 at intervals of 0.0001 and from 1 to 5.6 at intervals of 0.001.

(1941) XXVIII+302 pages; bound in buckram, \$2.00.

[Continued on p. 4 of cover]

[Continued from p. 3 of cover]

MT9. TABLE OF NATURAL LOGARITHMS, VOLUME II:

Logarithms of the integers from 50,000 to 100,000 to 16 places of decimals.
(1941) XVIII+501 pages; bound in buckram, \$2.00.

MT10. TABLE OF NATURAL LOGARITHMS, VOLUME III:

Logarithms of the decimal numbers from 0.0001 to 5.0000, to 16 places of decimals.
(1941) XVIII+501 pages; bound in buckram, \$2.00.

MT11. TABLES OF THE MOMENTS OF INERTIA AND SECTION MODULI OF ORDINARY ANGLES CHANNELS, AND BULB ANGLES WITH CERTAIN PLATE COMBINATIONS:

(1941) XIII+197 pages; bound in green cloth. \$2.00.

MT12. TABLE OF NATURAL LOGARITHMS, VOLUME IV:

Logarithms of the decimal numbers from 5.0000 to 10.0000, to 16 places of decimals.
(1941) XXII+506 pages; bound in buckram, \$2.00.

MT13. TABLE OF SINE AND COSINE INTEGRALS FOR ARGUMENTS FROM 10 TO 100:

(1942) XXXII+185 pages; bound in buckram, \$2.00.

MT14. TABLES OF PROBABILITY FUNCTIONS, VOLUME II:

Values of these functions to 15 places of decimals from 0 to 1 at intervals of 0.0001 and from 1 to 7.8 at intervals of 0.001.

(1942) XXI+344 pages; bound in buckram, \$2.00.

MT15. The hypergeometric and Legendre functions with applications to integral equations of potential theory. By Chester Snow, National Bureau of Standards. Reproduced from original handwritten manuscript.

(1942) VII + 319 pages; bound in heavy paper cover. \$2.00.

Payment is required in advance. Make remittance payable to the "National Bureau of Standards," and send with order, using the blank form facing page 3 of the cover.

The prices are for delivery in the United States and its possessions and in countries extending the franking privilege. To other countries the price of MT1 is 65 cents; that of MT2 to MT15, inclusive, is \$2.50 each; remittance to be made payable in United States currency.

Copies of these publications have been sent to various Government depositories throughout the country, such as public libraries in large cities, and colleges and universities, where they may be consulted.

A mailing list is maintained for those who desire to receive announcements regarding new tables as they become available.